

International 58th Meat Industry Conference “Meat Safety and Quality: Where it goes?”

Environmental performance of the poultry meat chain – LCA approach

Dubravka Skunca^{a,*}, Igor Tomasevic^b, Ilija Djekic^c

^aUniversity of Belgrade, Faculty of Agriculture, Nemanjina 6, Belgrade 11080, Serbia

^bAnimal Source Food Technology Department, University of Belgrade, Faculty of Agriculture, Nemanjina 6, Belgrade 11080, Serbia

^cDepartment of Food Safety and Quality Management, University of Belgrade, Faculty of Agriculture, Nemanjina 6, Belgrade 11080, Serbia

Abstract

This review aims to give an overview of published environmental assessments using the life cycle assessment (LCA) approach. LCA was deployed in terms of five main subsystems in the poultry meat chain: poultry farm, slaughterhouse, meat processing plant, retail and household use. This review revealed that 15 different environmental potentials are used as environmental indicators for estimating environmental performance of the poultry meat chain. General finding is that further research should use the LCA approach to assess the environmental performance of an overall poultry meat chain, focusing on the global warming potential, acidification potential, eutrophication potential and ozone layer depletion.

Keywords: life-cycle assessment; poultry meat chain; environmental potentials

1. Introduction

The environmental impact of livestock production has a major impact on the environment, since meat contributes between 4.6 and 7.1 gigatonnes of greenhouse gases (GHG) each year to the atmosphere and production processes for meat account for between 15% and 24% GHG emissions¹. The environmental impact of poultry chain is estimated to emit 0.6 gigatonnes CO₂-equivalent².

* Corresponding author. Tel: +381- 63- 841- 3194
E-mail address: dubravkaskunca@hotmail.com

According to ISO 14040, life-cycle assessment (LCA) represents the compilation and evaluation of the inputs, outputs and potential environmental impacts of a product system throughout its life cycle and it is a tool for the analysis of the environmental burden of products at all stages in their life cycle³. A limited number of studies have researched environmental performance in the meat chain, especially the poultry chain. There are papers targeting different aspects of the poultry meat chain, but there is an evident lack of studies concerning life-cycle assessment (LCA) approach for the environmental performance of an overall poultry meat chain. Table 1 gives an overview of the poultry chain LCA manuscripts emphasizing system boundaries and environmental impacts.

Table 1. Summary of studies linking environmental impacts of the poultry chain.

Authors	Sample	Research focus	System boundaries ^a					Environmental impact ^b
			1	2	3	4	5	
[4]	3 main broiler production systems in the UK	LCA of broiler production systems			✓			GWP, EP, AP, EC, LC, ADP
[5]	Two production scenarios of chickens	Comparison of production of chickens consumed in France	✓	✓	✓			GWP, CED, AP, EP, TEP, EC, LC
[6]	Chicken product	Environmental improvement through LCA methodology	✓	✓	✓	✓	✓	GWP, OLD, AP, EP, WC, CED
[7]	Chicken meat production systems in Switzerland	Environment impacts and selected import sources	✓				✓	CED, GWP, OLD, LC, AP, EP, TEP, HT, WC, FEP
[8]	Poultry slaughterhouse in Brazil	A case study of cleaner production		✓				WC, EP, FD
[9]	Two chicken meals	Life cycle assessment of integrated food chains – Sweden					✓	GWP, AP, EP, POFP
[10]	The US broiler poultry sector	Life cycle energy use and greenhouse gas, ozone depleting, acidifying and eutrophying emissions	✓	✓	✓			GWP, OLD, AP, EP, EC
[11]	Two differing Brazilian poultry production systems	Cradle to gate study	✓	✓	✓	✓		GWP, OLD, AP, EP
[12]	Finnish broiler chicken fillet product	Supply chain integrated LCA method	✓	✓	✓	✓		GWP, AP, EP, OLD, PS, FEP
[13]	Meat production	An estimation of potential future greenhouse gas emissions			✓			GWP

^a: Subsystem 1 — Poultry farm; Subsystem 2 – Slaughterhouse; Subsystem 3 – Meat processing plant; Subsystem 4 - Retail; Subsystem 5 - House hold use;

^b: Global warming potential (GWP); acidification potential (AP); eutrophication potential (EP), ozone layer depletion (OLD), photochemical smog (PS) and human toxicity (HT); abiotic depletion potential (ADP); land competition / use (LC); photochemical oxidants formation (POFP); energy consumption (EC); water consumption (WC); cumulative non-renewable fossil and nuclear energy demand (CED), terrestrial ecotoxicity (TEP), Freshwater depletion (FD), fresh water aquatic ecotoxicity (FEP)

2. Life-cycle assessment

LCA is a tool used for identifying hot spots in the production chain which may give opportunities for lowering environmental impacts while improving efficiency and profitability^{14,15}. Analysis of Table 1 shows that common potentials analyzed by all authors for subsystem 1 are GWP, AP and EP, for subsystem 2 is EP, for subsystem 3 is GWP, for subsystem 4 are GWP, AP, EP and OLD and for subsystem 5 are GWP, AP and EP.

The system boundaries cover five main subsystems ‘poultry farm’, ‘slaughterhouse’, ‘meat processing plant’, ‘retail’ and ‘household use’. The ‘waste and waste water’ subsystem is present in each of the five subsystems, which includes all activities related to waste management of solid waste and treatment of waste water.




Fig. 1. Generic model of system boundaries of the poultry meat production life cycle.

3. Discussion

Nine out of ten selected studies in Table 1 examined global warming potential (GWP) as important part of poultry meat chain. Research of prediction of environmental impacts of broiler production systems using LCA has found that processing and transport have large environmental impacts, especially in terms of the GWP⁴. Broiler chicken housing and related fodder production was responsible for most of the GWP, as production of fodder accounted for 36% of the total impact, and broiler housing 29% and this result was not only influenced by the emissions from energy consumption, but also by the nitrous oxide emissions from fertiliser production and use, as well as in nitrous oxide and methane emissions from handling broiler chicken manure¹². In the study concerning estimation of potential future GHG emissions from meat production it was found that meat production (beef, chicken and pig meat) will stay a large producer of greenhouse gases and it will account for up to 6.3% of current greenhouse gas emissions in 2030¹³.

In eight examined studies, acidification potential (AP) was regarded as an important environmental impact. As ammonia is an acidifying substance emitted from livestock production, the chicken farm is a large contributor to AP and the reduction in waste lowers acidifying emissions the most⁹. Pardo et al. concluded that up to 20% of GHG emissions connected to packaging were avoided by using biodegradable materials instead of plastic tray and film and that there was reduction in terrestrial acidification (-0.5%), as well as in ozone depletion (-1.5%)⁶. While researching the chicken meat supply chain and environmental impacts for its improvement, Katajajuuri et al. concluded that broiler housing had a large influence on acidification and eutrophication, because of nutrient run-off and leaching and ammonia emissions from broiler chicken manure¹².

Nine selected studies researched eutrophication potential (EP). Study of management of water use at a poultry slaughterhouse showed that dry cleaning procedures, control over the amount of water used on a sector-by-sector basis, better collection of exsanguination residues, automation of the water system and change in the configuration of the wastewater treatment plant through the use of an anaerobic reactor and a sequential oxic/anoxic system for reduction of the eutrophication effect of the final wastewater product should be adopted⁸.

Ozone layer depletion (OLD) has been examined in five studies. Production of fertilizers, and especially nitrogen fertilizer, was crucial for ozone depleting emissions¹⁰. By applying poultry litter to agricultural fields, litter

management offsets 1% of global warming emissions, 25% of ozone depleting emissions, 7% of acidifying emissions and 2,5% of eutrofying emissions¹⁰. Katajajuuri et al. concluded that broiler housing is the most important phase regarding tropospheric ozone formation because of the methane emissions from broiler manure. By using alternative fuels 70% of greenhouse gas emissions from broiler houses could be cut, but it would result in a 7% increase in tropospheric ozone formation because of increased air emissions¹².

4. Conclusion

Implementation of a LCA approach may lead to improvements of sustainability of the poultry meat chain. Further research should use the LCA approach for the environmental performance of an overall poultry meat chain including all the subsystems and focusing on the global warming potential (GWP), acidification potential (AP), eutrophication potential (EP) and ozone layer depletion (OLD). These four environmental potentials have been recognized as the most important in the poultry meat chain, but other environmental potentials should also be further investigated.

References

1. Steinfeld H, Gerber P, Wassenaar T, Castel V, Rosales M, Haan CD. Livestock's long shadow: environmental issues and options. *FAO*, Rome; 2006.
2. MacLeod M, Gerber P, Mottet A, Tempio G, Falcucci A, Opio C, Vellinga T, Henderson B, Steinfeld H. Greenhouse gas emissions from pig and chicken supply chains – A global life cycle assessment. In: *FAO*, Rome; 2013.
3. ISO 14040:2006. Environmental management — Life cycle assessment — Principles and framework, in *ISO*, Geneva, Switzerland.
4. Leinonen I, Williams A, Wiseman J, Guy J, Kyriazakis I. Predicting the environmental impacts of chicken systems in the United Kingdom through a life cycle assessment Broiler production systems. *Poultry Sci* 2012;**91**:8-25.
5. da Silva Junior VP, Cherubini E, Soares SR. Comparison of two production scenarios of chickens consumed in France. In: *8th International Conference on LCA in the Agri-Food Sector*, Saint-Malo, France; 2012. p. 546-51.
6. Pardo G, Ciruelos A, Lopez N, Gonzalez L, Ramos S, Zufia J. Environment improvement of a chicken product through life cycle assessment methodology. In: *8th Conference on LCA in the Agri-Food Sector*, Saint-Malo, France; 2012. p. 86-91.
7. Grandl F, Alig M, Mieleitner J, Nemecek T, Gaillard G. Environmental impacts of different pork and chicken meat production systems in Switzerland and selected import sources. In: *8th International Conference on LCA in the Agri-Food Sector*, INRA, Saint-Malo, France; 2012. p. 554-59.
8. Kist LT, El Moutaqi S, Machado ÊL. Cleaner production in the management of water use at a poultry slaughterhouse of Vale do Taquari, Brazil: a case study. *J Clean Prod* 2009;**17**:1200-5.
9. Davis J, Sonesson U. Life cycle assessment of integrated food chains—a Swedish case study of two chicken meals. *Int J Life Cycle Ass* 2008;**13**:574-84.
10. Pelletier N. Environmental performance in the US broiler poultry sector: Life cycle energy use and greenhouse gas, ozone depleting, acidifying and eutrophying emissions. *Agr Syst* 2008;**98**:67-73.
11. da Silva Júnior VP, Soares S, de Alvarenga R. Cradle to gate study of two differing Brazilian poultry production systems. In: *6th International Conference on LCA in the Agri-Food Sector*. Zurich; 2008.
12. Katajajuuri JM, Grönroos J, Usva K. Environmental impacts and related options for improving the chicken meat supply chain. In: *6th International Conference on LCA in the Agri-Food Sector*. Zurich, 2008.
13. Fiala N. Meeting the demand: An estimation of potential future greenhouse gas emissions from meat production. *Ecol Econ* 2008;**67**:412-19.
14. Djekic I, Miocinovic J, Tomasevic I, Smigic N, Tomic N. Environmental life-cycle assessment of various dairy products. *J Clean Prod* 2014;**68**:64-72.
15. Eide MH. Life cycle assessment (LCA) of industrial milk production. *Int J Life Cycle Ass* 2002;**7**:115-26.